



Testing inter-teeth variability in adult individual age-at death estimate using cementochronology (TCA)

Laëtitia Lanteri, Aurore Schmitt, Bruno Foti, Stephan Naji

► To cite this version:

Laëtitia Lanteri, Aurore Schmitt, Bruno Foti, Stephan Naji. Testing inter-teeth variability in adult individual age-at death estimate using cementochronology (TCA). 82nd Annual Meeting of the American Association of Physical Anthropologists, Apr 2013, Knoxville, United States. 150, pp.177-177, 2013. halshs-01074756

HAL Id: halshs-01074756

<https://shs.hal.science/halshs-01074756>

Submitted on 8 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Testing inter-teeth variability in adult individual age-at-death estimate using cementochronology (TCA)

Laëticia LANTERI, Aurore SCHMITT, Bruno FOTI and Stephan NAJI


Laboratoire d'Anthropologie Bioculturelle, Aix-Marseille Université
Bd Pierre Dramard, 13344 Marseille cedex 15, France

Correspondance to Laëticia Lanteri: laetitia.lanteri@etu.univ-amu.fr

Introduction

Do all teeth of one individual indicate the same age?

- Variability of cementum only mentioned by 2 studies:
 - Charles *et al.* (1986): 43% between canines and premolars
 - Wittwer-Backofen *et al.* (2004): "no consistent high error"



Objectives

Studying intra-individual variability of the dental cementum

- Repeatability and reproducibility of counts
- Between dental classes
- Between teeth of one individual

Materials and Methods


- 51 modern single-rooted teeth
- 15 individuals (39 – 81 years old)
- Central and lateral incisors, canines and premolars
- Multiple extractions from 2 to 12 teeth per individual

Protocol:

Preparation

- Anonymisation of teeth and sections
- Embedding teeth in epoxy resin

Sectioning



Counting

- Manual
- Area with the **largest number** of lines
- Three counts** per tooth minimum

Statistical tests:

- Intra and inter-observer variability
- Link between error and tooth class (Kruskal-Wallis test + paired comparison)
- Average and maximum number of counted lines

Results

- 11 teeth excluded
 - Fragility
 - Unreadable
 - Insufficient number of counts (< 3/5)
- 4 individuals represented by 1 tooth



Intra and inter observer variability

| Intra observer | Inter observer |
|----------------|----------------|
| 0.76 | 0.60 |

No link between error and class of teeth ($p > 0.05$)

Between teeth of one individual

Sample: 36 teeth of 10 individuals

- Using maximum of the age estimations:


- Average deviation per individual:

2 – 13 years

- Minimal and maximal deviation per individual:

0 – 26 years

- Entire sample: **6.5 years** of average deviation



Between dental classes

Sample: 40 teeth of 14 individuals

| Average error | |
|--------------------------------|--------------------------------|
| Maximum of the age estimations | Average of the age estimations |
| 2.64 years | 8.43 years |

- Incisors ($n=33$) – lowest errors with:
 - Maximum of the estimations
 - Mandibular central incisors

Discussion

- Repeatability and reproducibility of counts: **"good" coefficients (0.76 and 0.60)**
- Maximal** estimation of age is more accurate than the average one
- Between dental classes:
 - Age structure: overestimation of age of younger adults and underestimation of age of older adults
 - Mandibular central incisors** : the lowest error
- Between teeth of one individual:
 - Average deviation: **6.5 years** (acceptable; similar to other methods)
 - Up to 26 years** (premolars known to be problematic)

Conclusion

- Proved presence of intra individual variability between dental classes **AND** between teeth of one individual, sometimes very high (26) protocol to improve further
- Maximal number of incremental lines provides more accurate and reliable estimations of age
- Prospects: greater number of teeth, intra tooth variability, sexual and pathological influences